

On generalized topological spaces

ARTUR PIĘKOSZ

Abstract

In this paper a systematic study of the category **GTS** of generalized topological spaces (in the sense of H. Delfs and M. Knebusch) and their strictly continuous mappings begins. It is then applied in reconstructing the important elements of the theory of locally definable and weakly definable spaces over (weakly) topological structures.

2000 MS Classification: 54A05, 18F10, 03C65.

Key words and frases: generalized topological space, Grothendieck topology, topological structure.

1 Introduction

In 1991, in his paper [7], M. Knebusch suggested that his theory of locally semialgebraic (developed in [3] together with H. Delfs) and weakly semialgebraic (developed in [6]) spaces could be generalized to the o-minimal context. This was not done for many years until recently (see [10]), but the theory is still not broadly understood. Especially the concept of a *generalized topological space*, defined on page 1 of [3], seems to make a problem. (Notice that the paper [1] does not mention generalized topology at all. Moreover, the considerations in [1] use in practice the non-fact that generalized topology reduces to topology.) This paper aims to solve this problem by looking closer at generalized topological spaces.

One way to help in this situation is to generalize the theory to see which elements play which roles. The category **GTS** of generalized topological spaces and their strictly continuous mappings (probably the most important concept of the mentioned volumes of Lecture Notes in Mathematics) needs more explanation, as it is an alternative to the usually used category **Top** of topological spaces and continuous mappings. It originates from the categorical concept of Grothendieck topology. **GTS** contains **Top** as a full subcategory. Only the use of **GTS** allowed to glue infinitely many definable sets to produce locally definable spaces in [3] and [10] and weakly definable spaces in [6] and [10]. Many of the proofs of [3] and [6] are purely topological (naturally in the sense of the generalized topology), and it seems to be important to extract the information on particular levels of structure involved in the construction of a locally or weakly definable space. It appears that then the theory of infinite gluings of definable sets can be reconstructed to a large extent in a more general setting of weakly topological structures (maybe with some additional assumptions). The author is unaware of any regular study of the category **GTS** on its own, thus decided to make the first step.

This paper is a continuation of [10]. It gives a basic theory of the category **GTS**. In particular, the paper deals with main (generalized) topological concepts as: small sets, bases, connectedness, completeness, paracompactness, Lindelöfness, separation axioms. Existence of colimits and products are considered. Many examples are given. The reader may notice the diversity of discrete spaces in **GTS** and their important role in the proofs. Then **GTS** is used to build categories of locally definable and weakly definable spaces

over model-theoretic structures. The minimal setting for topological considerations is assuming a topology on the underlying set M of a model-theoretic structure $\mathcal{M}$, and insisting on regarding the product topologies on the cartesian powers of M . Such structures are called in this paper weakly topological structures. They are more general than so called first order topological structures of [11, 9] and structures with a definable topology of [12].

2 Grothendieck topology

Here we remind what a Grothendieck topology is. The reader may consult books like [2, 5, 8].

Let $\mathcal{C}$ be a small category. Consider the category of presheaves of sets on $\mathcal{C}$, denoted by $Psh(\mathcal{C})$ or $\hat{\mathcal{C}}$, which is the category of contravariant functors from $\mathcal{C}$ to **Sets**. Let us remind the fundamental fact:

Fact 2.1 (Yoneda lemma, weak version). *Functor $\mathcal{C} \ni C \mapsto Hom(-, C) \in \hat{\mathcal{C}}$ is full and faithful, so we can consider $\mathcal{C} \subseteq \hat{\mathcal{C}}$.*

To define a Grothendieck topology the following notion is usually used: A **sieve** S on an object C is a subobject of $Hom(-, C) \in \hat{\mathcal{C}}$. Since a sieve on C is a presheaf of sets of morphisms with common codomain C , this may be translated (with a small abuse of language) into: S is a set of morphisms with codomain C such that $f \in S$ implies $f \circ g \in S$, if only $f \circ g$ has sense. The largest sieve on C is $Hom(-, C)$, the smallest is $\emptyset$.

A **Grothendieck topology** J on $\mathcal{C}$ is a function $C \mapsto J(C)$, with $J(C)$ a set of sieves on C such that the following axioms hold:

(**identity/nonemptiness**) for each C , $Hom(-, C) \in J(C)$;

(**stability/base change**) if $S \in J(C)$ and $f : D \rightarrow C$ is a morphism of $\mathcal{C}$, then $f^*S = \{g \mid f \circ g \in S\} \in J(D)$;

(**transitivity/local character**) if $S \in J(C)$ and R is a sieve on C such that $f^*R \in J(D)$ for each $f \in S$, $f : D \rightarrow C$, then $R \in J(C)$.

Elements of $J(C)$ are called **covering sieves**. Pair $(\mathcal{C}, J)$ is called a **Grothendieck site**. The above axioms imply the conditions (see section III.2 of [8]):

(**saturation**) if $S \in J(C)$ and R is a sieve containing S , then $R \in J(C)$;

(**intersection**) if $R, S \in J(C)$, then $R \cap S \in J(C)$.

It follows that each $J(C)$ is a filter (not necessary proper) on the lattice $Sub_{\hat{\mathcal{C}}}Hom(-, C)$ of sieves on C .

If the category $\mathcal{C}$ has *pullbacks*, then instead of covering sieves, we can speak about **covering families** of morphisms (generating respective sieves), so the axioms may be rewritten as:

(**identity/isomorphism**) for each C , $\{id_C\}$ is a covering family (stated also: for each isomorphism $f : D \rightarrow C$, $\{f\}$ is a covering family);

(**base change**) if $\{f_i : U_i \rightarrow U\}_i$ is a covering family, and $g : W \rightarrow U$ any morphism, then $\{\pi_{2i} : U_i \times_U W \rightarrow W\}_i$ is a covering family;

(**local character**) if $\{f_i : U_i \rightarrow U\}_i$ is a covering family, and $\{g_{ij} : V_{ij} \rightarrow U_i\}_j$ are covering families, then $\{f_i \circ g_{ij} : V_{ij} \rightarrow U\}_{ij}$ is a covering family;

and usually the following is added (see Definition 16.1.2 of [5]):

(**saturation**) if $\{f_i : U_i \rightarrow U\}_i$ is a covering family, and each of the f_i 's factorizes through an element of $\{g_j : V_j \rightarrow U\}_j$, then $\{g_j : V_j \rightarrow U\}_j$ is a covering family.

Alternatively (see section 6.7 of [2]) authors consider saturated and non-saturated Grothendieck topologies, but for a Grothendieck site saturation is usually assumed.

Grothendieck topology allows to define sheaves (of sets). A **sheaf** is such a presheaf F that for each covering family $\{U_i \rightarrow U\}_i$ in the respective diagram

$$F(U) \xrightarrow{e} \prod_i F(U_i) \xrightarrow[p_2]{p_1} \prod_{i,j} F(U_i \times_U U_j)$$

the induced morphism e is the equalizer of the standardly considered pair of morphisms p_1, p_2 (cf. section III.4 in [8]).

A Grothendieck topology is **subcanonical** if every representable presheaf is a sheaf, so in this case we may, by identifying the objects of $\mathcal{C}$ with their respective representable presheaves, consider $\mathcal{C} \subseteq Sh(\mathcal{C}) \subseteq Psh(\mathcal{C}) = \hat{\mathcal{C}}$. Grothendieck topologies used in practice are usually subcanonical.

3 Generalized topological spaces

Now, for any set X , we have a boolean algebra $\mathcal{P}(X)$ of subsets of X , so it may be treated as a small category with inclusions as morphisms. In this category fibered products are the same as (binary) products and the same as (binary) intersections, so $\mathcal{P}(X)$ has pullbacks. We want to introduce a full subcategory of $\mathcal{P}(X)$, consisting of “open subsets” of X . Then we want to introduce a subcanonical Grothendieck topology on this category. Subcanonicity means in this setting that for each covering family of morphisms (which may be identified with a family of subsets of a given set, since morphisms are inclusions) the object covered by the family is the supremum of this family in the smaller category.

This leads to the notion of a generalized topological space introduced by H. Delfs and M. Knebusch in [3].

A **generalized topological space** (gts) is a set X together with a family of subsets $\overset{\circ}{\mathcal{T}}(X)$ of X , called **open sets**, and a family of open families Cov_X , called **admissible families** or **admissible (open coverings)**, such that:

- (A1) $\emptyset, X \in \overset{\circ}{\mathcal{T}}(X)$ (the empty set and the whole space are open),
- (A2) if $U_1, U_2 \in \overset{\circ}{\mathcal{T}}(X)$ then $U_1 \cup U_2, U_1 \cap U_2 \in \overset{\circ}{\mathcal{T}}(X)$ (finite unions and finite intersections of open sets are open),
- (A3) if $\{U_i\}_{i \in I} \subset \overset{\circ}{\mathcal{T}}(X)$ and I is finite, then $\{U_i\}_{i \in I} \in \text{Cov}_X$ (finite families of open sets are admissible),

The above three axioms are a strengthening of the identity axiom. They also insure that the smaller category has pullbacks. These three axioms may be collectively called the **finiteness** axiom.

- (A4) if $\{U_i\}_{i \in I} \in \text{Cov}_X$ then $\bigcup_{i \in I} U_i \in \overset{\circ}{\mathcal{T}}(X)$ (the union of an admissible family is open),

This axiom may be called **co-subcanonicity**. Together with subcanonicity, it means that admissible families are coverings (in the traditional sense) of their unions, which is imposed by the notation of [3]:

$\{U_i\}_{i \in I} \in \text{Cov}_X(U)$ iff U is the union of $\{U_i\}_{i \in I}$.

Subcanonicity and co-subcanonicity may be collectively called the **naturality** axiom.

- (A5) if $\{U_i\}_{i \in I} \in \text{Cov}_X$, $V \subset \bigcup_{i \in I} U_i$, and $V \in \mathring{\mathcal{T}}(X)$, then $\{V \cap U_i\}_{i \in I} \in \text{Cov}_X$ (the intersections of an admissible family with an open subset of the union of the family form an admissible family),

This is the stability axiom.

- (A6) if $\{U_i\}_{i \in I} \in \text{Cov}_X$ and for each $i \in I$ there is $\{V_{ij}\}_{j \in J_i} \in \text{Cov}_X$ such that $\bigcup_{j \in J_i} V_{ij} = U_i$, then $\{V_{ij}\}_{\substack{i \in I \\ j \in J_i}} \in \text{Cov}_X$ (all members of admissible coverings of members of an admissible family form together an admissible family),

This is the transitivity axiom.

- (A7) if $\{U_i\}_{i \in I} \subset \mathring{\mathcal{T}}(X)$, $\{V_j\}_{j \in J} \in \text{Cov}_X$, $\bigcup_{j \in J} V_j = \bigcup_{i \in I} U_i$, and $\forall j \in J \exists i \in I : V_j \subset U_i$, then $\{U_i\}_{i \in I} \in \text{Cov}_X$ (a coarsening, with the same union, of an admissible family is admissible),

This is the saturation axiom.

- (A8) if $\{U_i\}_{i \in I} \in \text{Cov}_X$, $V \subset \bigcup_{i \in I} U_i$ and $V \cap U_i \in \mathring{\mathcal{T}}(X)$ for each i , then $V \in \mathring{\mathcal{T}}(X)$ (if a subset of the union of an admissible family forms open intersections with the members of the family then the subset is open). This axiom may be called **regularity**.

Both saturation and regularity have a smoothing character. Saturation may be achieved by adding coarsenings of admissible coverings for any *gts*. Regularity may be achieved for the class of locally small spaces (see below for the definition) by adding “locally open” subsets and allowing locally essentially finite coverings (see section I.1 in [3]).

A **strictly continuous mapping** between *gtses* is such a mapping that the preimage of an open set is open and the preimage of an admissible covering is an admissible covering. (They may be considered morphisms of sites in this context, compare section 6.7 of [2] and section 17.2 of [5].) The *gtses* together with the strictly continuous mappings form a category called here **GTS**. Isomorphisms of **GTS** will be called **strict homeomorphisms**.

We call a subset K of a *gts* X **small** if for each admissible covering $\mathcal{U}$ of any open U , the set $K \cap U$ has a finite subcovering. (We say in this case that the covering $\mathcal{U}$ is **essentially finite** on K or on $K \cap U$.) The class of small spaces forms a full subcategory **SS** of **GTS**.

For families of sets $\mathcal{U}, \mathcal{V}$, we will use usual set-theoretic operations $\cup, \cap, \setminus$ on sets, and operations on families of sets, for example

$$\mathcal{U} \sqcup \mathcal{V} = \{U \cup V : U \in \mathcal{U}, V \in \mathcal{V}\}, \quad \mathcal{U} \times \mathcal{V} = \{U \times V : U \in \mathcal{U}, V \in \mathcal{V}\}.$$

In particular, we will denote $V \sqcup \mathcal{U} = \{V\} \sqcup \mathcal{U}$ for a set V . Notice that a subfamily of an admissible family may not be admissible, even if they have the same union. Similarly if $\mathcal{U}$ and $\mathcal{V}$ are admissible, then $\mathcal{U} \cap \mathcal{V}$ may not be admissible, but $\mathcal{U} \sqcup \mathcal{V}$ is always admissible

by stability and transitivity. On the other hand, $\mathcal{U} \cup \mathcal{V}$ is always admissible for admissible $\mathcal{U}, \mathcal{V}$ by finiteness and transitivity.

Theorem 3.1. 1) A subset of a small set is small.

2) The image of a small set by a strictly continuous mapping is small.

Proof. 1) Take any admissible (open) covering $\mathcal{U}$ of U . Assume $L \subseteq K$ and K is small. Since $\mathcal{U}$ is essentially finite on $K \cap U$, it is also essentially finite on its subset $L \cap U$.

2) Assume $f : X \rightarrow Y$ is strictly continuous, and $K \subseteq X$ is small. Take an admissible open covering $\mathcal{V}$ of V in Y . Then $f^{-1}(\mathcal{V})$ is essentially finite on K , and $f(K) \sqsubseteq \mathcal{V} = f(K \sqcap f^{-1}(\mathcal{V}))$ is essentially finite. $\square$

Theorem 3.2. In any gts if an open family is essentially finite (on its union), then this family is admissible.

Proof. First notice that if an open family $\mathcal{U}$ has the largest element U , then $\{U\}$ and $\mathcal{U}$ are refinements of each other. Thus, by saturation, $\mathcal{U}$ is admissible.

Now if an open family $\mathcal{U}$ has a finite subcover $\mathcal{U}_0$ (of its union), then for each $V \in \mathcal{U}_0$ the family $V \sqsubseteq \mathcal{U}$ has the largest element V , and is admissible. Since $\mathcal{U}_0$ is admissible, also $\mathcal{U}_0 \sqsubseteq \mathcal{U}$ is admissible by transitivity, and $\mathcal{U}$ is admissible by saturation. $\square$

Corollary 3.3. An open family of a small space is admissible if and only if it is essentially finite.

Example 3.4. Each topological space may be considered as a gts. Admissible coverings are understood as any coverings in the traditional sense. Each continuous function is here strictly continuous. Thus **Top** is a full subcategory of **GTS**.

Proposition 3.5. The real line with the usual topology (denoted $\mathbb{R}_{top}$) is a gts in which all small subsets are finite.

Proof. A small subset of the (topological) space $\mathbb{R}_{top}$ has only finitely many connected components. Each such component is an interval of some type. Each open subset of such an interval must be quasi-compact, so the interval must be degenerated to a point. Thus any small subset is finite. $\square$

Example 3.6. Given a topological space, we could consider only singletons of open sets as admissible coverings of these sets. This would give so-called **indiscrete Grothendieck topology**, but not (in general) a gts, because of (A3). On the other hand, if we considered all families of open subsets as coverings of an open set, the resulting **discrete Grothendieck topology** would not (in general) be subcanonical, hence not a gts.

Example 3.7 (“the subanalytic site,” Remark 21 in [10]). If M is a real analytic manifold, then we can set:

- a) open subset of a gts = open subanalytic subset;
- b) admissible covering of a gts = covering that is essentially finite on compact subsets.

Example 3.8 (another “site,” Remark 21 in [10]). In the situation as in Example 3.7, set:

- a) open subset of a gts = any subanalytic subset;
- b) admissible covering of a gts = covering that is essentially finite on compact subsets.

Notice that compact sets are small in Examples 3.7 and 3.8, but not necessarily small in Example 3.4 (because an open subset of a compact set is not usually compact).

For any *gts* X and a small or open $Y \subseteq X$ we can induce a *gts* on Y (then Y will be called a **subspace** of X) in the following way:

- a) open subsets of the *gts* Y = traces of open subsets of the *gts* X on Y ,
- b) admissible families of the *gts* Y = traces of admissible families of the *gts* X on Y .

This works for open subsets, since openness and admissibility in an open subspace is equivalent to openness and admissibility in the whole space. It works for small subsets because in a small space “admissible” means exactly “essentially finite”. In general, the problem with transitivity, saturation and regularity arises.

A subset Y of a *gts* X is **closed** if its complement Y^c is open, and is **locally closed** if it is an intersection of a closed set and an open set. A subset Y of a *gts* X is **constructible** if it is a Boolean combination of open sets. Remind that each constructible set is a finite union of locally closed sets.

If $\{(\overset{\circ}{\mathcal{T}}_\alpha, Cov_\alpha)\}_{\alpha \in A}$ is a family of generalized topologies on a set X , then their intersection $(\bigcap_\alpha \overset{\circ}{\mathcal{T}}_\alpha, \bigcap_\alpha Cov_\alpha)$ is a generalized topology. Thus, for any family $\{V_i\}_{i \in I}$ of subsets of a *gts* X and any family $\{\mathcal{U}_j\}_{j \in J}$ of subfamilies of $\{V_i\}_{i \in I}$ (this is not restrictive because we may enlarge this family if needed), we can speak about the generalized topology **generated by** $\{V_i\}_{i \in I}$ and $\{\mathcal{U}_j\}_{j \in J}$. A **basis** of the generalized topology is such a family of open sets that any open set is an admissible union of elements from the basis. (The notion of the basis of a topology is a special case of the notion of the basis of the generalized topology.) Notice that if $\mathcal{B}$ is a basis of a *gts* $(X, \overset{\circ}{\mathcal{T}}, Cov)$, then $(\mathcal{B}, Cov^{\mathcal{B}})$ generates $(\overset{\circ}{\mathcal{T}}, Cov)$, where $Cov^{\mathcal{B}} = Cov \cap \mathcal{P}(\mathcal{B})$.

A subset $O \subseteq X$ will be called **weakly open** if it is a union of open subsets of the *gts* X , and **weakly closed** if its complement is weakly open. The weakly open subsets form a topology generated by the open subsets from the generalized topology. (For the class of locally small spaces the generated topology will be called the strong topology.) Notice that a basis of a generalized topology is also a basis of its generated topology.

The following example shows that a continuous mapping (in the generated topology) that maps small sets onto small sets may not be strictly continuous.

Example 3.9. Take the real line $\mathbb{R}$ and define the open sets as the finite unions of open intervals with endpoints being rational numbers or infinities. Define the admissible coverings as the essentially finite coverings. We get a small *gts*. The mapping $\mathbb{R} \ni x \mapsto rx \in \mathbb{R}$, for $r \notin \mathbb{Q}$, is continuous but not strictly continuous.

A subset $Y \subseteq X$ will be called **dense** in X if its weak closure (i.e. closure in the generated topology) is equal to X . A *gts* X is **separable** if there is a countable dense subset of X .

We can introduce the countability axioms: a *gts* X satisfies the **first axiom of countability** if each point of X has a countable basis of open neighborhoods (such a family may easily be made admissible), and satisfies the **second axiom of countability** if there is a countable basis of the generalized topology. The semialgebraic real line $\mathbb{R}$ has a countable basis of the generated topology $\{(p, q) : p < q, p, q \in \mathbb{Q}\}$ which is not a basis of the generalized topology. Of course, each *gts* satisfying the second axiom of countability is separable.

The separation axioms in **GTS** have weak and strong versions. A *gts* X will be called: a) **weakly** T_1 if for each $x \in X$ and $y \in X \setminus \{x\}$ there is an open set U such that $x \in U$ and $y \notin U$; b) **strongly** T_1 if each singleton is a closed subset of X ; c) **weakly Hausdorff** if for each pair of different points $x, y \in X$ there are open disjoint U, V such that $x \in U$ and $y \in V$; d) **strongly Hausdorff** if it is weakly Hausdorff and strongly T_1 ; e) **weakly regular** if for each point $x \in X$ and each subset F of X not containing x and being either closed or a singleton there are open disjoint sets U, V such that $x \in U$ and $F \subseteq V$; f) **strongly regular** if it is strongly T_1 and weakly regular; g) **weakly normal** if for two disjoint sets F, G being each either closed or a singleton there are open disjoint sets U, V such that $F \subseteq U$ and $G \subseteq V$; h) **strongly normal** if it is strongly T_1 and weakly normal.

It is clear that a *gts* is weakly T_1 (weakly Hausdorff) if and only if the generated topology of X is T_1 (Hausdorff, respectively). Each weakly regular *gts* has a regular generated topology.

Example 3.10. *Let X be an infinite gts where open sets are exactly finite sets or the whole space, and admissible coverings are exactly essentially finite open families. Then X is not strongly T_1 , but the generated topology is discrete.*

Proposition 3.11. *The category **SS** has products.*

Proof. The product of a set of small spaces $\{X_\alpha\}_{\alpha \in A}$ is the cartesian product of sets $X = \prod_{\alpha \in A} X_\alpha$ with the generalized topology described as follows: open sets are finite unions of full cylinders with bases being finite cartesian products of open sets and admissible coverings are essentially finite coverings. Then obviously the projections $\pi_\alpha : X \rightarrow X_\alpha$ are strictly continuous, and for a system of strictly continuous mappings $f_\alpha : Y \rightarrow X_\alpha$, the induced mapping $(f_\alpha)_\alpha : Y \rightarrow X$ is strictly continuous. $\square$

An **open mapping** is a strictly continuous mapping between *gtss* such that the image of any open subset of the domain is an open subset of the range. It is clear that the canonical projections from the product of small spaces to its factors are open mappings. Similarly, a **closed mapping** maps closed subsets onto closed subsets. The canonical projections from the product of small spaces to its factors are closed mappings. Each strict homeomorphism is both an open mapping and a closed mapping. A **local strict homeomorphism** is a strictly continuous mapping $f : X \rightarrow Y$ for which there is an open admissible covering $X = \bigcup_{\alpha \in A} X_\alpha$ such that each $f|_{X_\alpha}$ is an open mapping and a strict homeomorphism onto its image. The next example shows that a local strict homeomorphism may not be an open mapping.

Example 3.12. *A locally semialgebraic covering $p : \mathbb{R}_{loc} \rightarrow S^1$ (see Example 11 in [10]) is a local strict homeomorphism but it is neither an open mapping nor a closed mapping. After passing to the strong topologies, we get the mapping $p_{top} : (\mathbb{R}_{loc})_{top} \rightarrow (S^1)_{top}$, which is open but not closed.*

Fact 3.13 (I.2 (p. 11) in [3]). *The category **GTS** has colimits.*

For any family $\{X_\alpha\}_{\alpha \in A}$ of *gtss* such that each intersection $X_\alpha \cap X_\beta$ is an open subspace both in X_α and in X_β , there is a unique *gts* X having all X_α s as open subspaces and the family $\{X_\alpha\}_{\alpha \in A}$ admissible. This is clear for finite families, and follows from the Fact 3.13 for any families, since then X is the colimit of the system of finite unions

of spaces X_α partially ordered by inclusion. In particular, if there is an admissible covering $\mathcal{U}$ of X , then the space X is uniquely determined as the admissible union of the family $\mathcal{U}$ of open subspaces. In such a situation, we will write $X = \bigcup^a \mathcal{U}$. If the members of $\mathcal{U} = \{U_i\}_{i \in I}$ are pairwise disjoint, then the resulting space X is called the **direct (generalized topological) sum** (or coproduct) of the elements of $\mathcal{U}$, and will be denoted $X = \bigoplus_{i \in I} U_i$. Notice that then each U_i is also closed. Moreover, each family of unions of U_i s is open (by regularity) and admissible (by saturation).

A *gts* X will be called **weakly discrete** if all its singltons are open subsets, **discrete** if all its subsets are open, and **topological discrete** if all families of subsets of X are open and admissible. (For a topological discrete space X , the formula $X = \bigoplus_{x \in X} \{x\}$ applies. All small topological discrete spaces are finite.) The space from Example 3.10 is weakly discrete but not discrete.

Example 3.14. *The semialgebraic space $\mathbb{R}$ has a small infinite weakly discrete subspace $\mathbb{N}$. The subspace $\mathbb{N}$ differs from the space from Example 3.10.*

Example 3.15 (infinite discrete small spaces). *On any infinite set X , there is still a generalized topology making X a discrete small space. It is enough to set: open subset is any subset, admissible family is any essentially finite family.*

Example 3.16. *The identity mapping from the infinite discrete small space on some (infinite) set X to the topological discrete space on this set is a closed and open strictly continuous bijection, but not a strict homeomorphism.*

A *gts* X will be called **connected** if X there is no pair U, V of open, disjoint, nonempty subsets of X such that $U \cup V = X$.

Proposition 3.17. *If X is a small space, $x \in X$, and $\{C_\alpha\}_{\alpha \in A}$ is a family of connected subsets of X each containing x , then $\bigcup_{\alpha \in A} C_\alpha$ is connected.*

Proof. Assume that the open subsets U, V of $\bigcup_{\alpha \in A} C_\alpha$ are disjoint and nonempty, cover $\bigcup_{\alpha \in A} C_\alpha$, and $x \in U$. Let $y \in V$. Then $y \in C_{\alpha_0}$ for some $\alpha_0 \in A$. The sets $U \cap C_{\alpha_0}$ and $V \cap C_{\alpha_0}$ are open, nonempty, and cover C_{α_0} , thus C_{α_0} is not connected, contradiction. Hence $\bigcup_{\alpha \in A} C_\alpha$ is connected. $\square$

The **connected component** of a point $x \in X$ of a small space X is the largest connected set C_x containing x . Since the weak closure of C_x is connected, C_x is weakly closed. The **quasi-component** $\tilde{C}_x$ of x is the intersection of clopen subsets of X containing x . Each quasi-component is also weakly closed. Each connected component is contained in a quasi-component.

We will say that a space X **satisfies** (ACC) if the family of connected components of X is open and admissible.

Proposition 3.18. *Each small space satisfying (ACC) has a finite number of connected components.*

Proof. Choose one point in any connected component of the space. The resulting subspace is topological discrete and small, so it is finite. $\square$

4 Locally small spaces

A *gts* is **locally small** if there is an admissible covering of the space by small open subsets. Locally small *gtses* form a full subcategory **LSS** of **GTS**. Notice that in a locally small *gts* each small set is contained in a small open set. A family of subsets of a locally small space is called **locally finite** if each open small set meets only finitely many members of the family.

Example 4.1. *Each topological discrete space is locally small (and it is not small, if infinite).*

Theorem 4.2. *An open family of a locally small space is admissible if and only if it is “locally essentially finite”, which means: the family is essentially finite on each small open subset.*

Proof. Each admissible family is locally essentially finite by the definition of a small subset. If an open family $\mathcal{V}$ is locally essentially finite, then it is essentially finite on members of an admissible covering $\mathcal{U}$ of the space by open small subspaces. It means that for each member U of $\mathcal{U}$, the family $U \sqcap \mathcal{V}$ is admissible. By the transitivity axiom, the family $\mathcal{U} \sqcap \mathcal{V}$ is admissible. By the saturation axiom, the family $\mathcal{V}$ is admissible. $\square$

Corollary 4.3. *Each locally finite open family in a locally small space is admissible.*

Theorem 4.4. *Each locally essentially finite union of closed subsets of a locally small space is closed.*

Proof. Let Z be a locally essentially finite union of closed sets. Take an admissible covering $\mathcal{U}$ of the space by small open subsets. For each element U of $\mathcal{U}$, the set $Z \cap U$ is a finite union of relatively closed subsets of U , so it is relatively closed. Each relative complement $Z^c \cap U$ is open in U . By regularity, the set Z^c is open. $\square$

Theorem 4.5. *The category **LSS** has finite products.*

Proof. For a family $X_1, \dots, X_k$ of locally small spaces assume that their admissible coverings by small open subsets $\mathcal{U}_1, \dots, \mathcal{U}_k$, respectively, are given. Then the product space is given by the formula

$$X_1 \times \dots \times X_k = \bigcup^a \mathcal{U}_1 \times \dots \times \mathcal{U}_k.$$

The projections $\pi_i : X_1 \times \dots \times X_k \rightarrow X_i$ are obviously strictly continuous, and for given strictly continuous $f_i : Y \rightarrow X_i$, the mapping $(f_1, \dots, f_k) : Y \rightarrow X_1 \times \dots \times X_k$ is also strictly continuous, since “admissible” means “locally essentially finite”. $\square$

The next example shows that the canonical projections along locally small spaces may not be open.

Example 4.6. *Consider the projection $\pi_2 : \mathbb{R}_{loc} \times \mathbb{R} \rightarrow \mathbb{R}$ (as locally semialgebraic spaces, see [3] or [10]). The image of an open locally semialgebraic set $\bigcup_{n \in \mathbb{N}} (n, n+1) \times (n, n+1)$ under π_2 is not an open subspace of $\mathbb{R}$.*

Fact 4.7. *If $\mathcal{U} = \{U_i\}_{i \in I}$ is a locally finite open family in some locally small space, and for each $i \in I$ some open $V_i \subseteq U_i$ is given, then $\{V_i\}_{i \in I}$ is locally finite, thus admissible.*

Fact 4.8. *If X is a small space and Y is a locally small space, then the projection $X \times Y \rightarrow Y$ is an open and closed mapping.*

On locally small spaces we introduce a topology, called the **strong topology**, whose basis is the family of open sets of the *gts*. (The members of the strong topology are exactly weakly open subsets of the space.) Passing to the strong topology gives a functor $(\)_{top} : \mathbf{LSS} \rightarrow \mathbf{Top}$.

A locally small space is called: **paracompact** if there is a locally finite covering of the space by small open subsets, and **Lindelöf** if there is a countable admissible covering of the space by small open subsets. Each connected paracompact locally small space is Lindelöf (the proof of I.4.17 in [3] is purely topological).

Theorem 4.9 (cf. I.4.6 of [3]). *For each paracompact locally small space X , the weak closure $\overline{Y}$ (that is: in the strong topology) of a small set Y is small.*

Proof. Take a locally finite covering $\mathcal{U}$ of X by small open subsets. The set Y is covered by a finite subcover $\mathcal{U}_0$ of $\mathcal{U}$ of all members of $\mathcal{U}$ that meet Y . Then $\overline{Y}$ and the union of $\mathcal{U} \setminus \mathcal{U}_0$ are disjoint, and $\overline{Y}$ is contained in the union of $\mathcal{U}_0$, which is a small set. $\square$

Example 4.10. *If a metric space (X, d) satisfies the ball property*

(BP) each intersection of two open balls is a finite union of open balls,

then X has a natural generalized topology, where an open set Y is such a subset of X that the trace of Y on each open ball is a finite union of open balls, and the admissible coverings are such open coverings that are essentially finite on each open ball. Then open balls are small sets. The covering of the space by all open balls is admissible (each small set is covered by one open ball), and X is a locally small space satisfying the first axiom of countability. The family of all open balls is a basis of the generalized topology.

A subset Y of a locally small space X is **locally constructible** if each intersection $Y \cap U$ with a small open $U \subseteq X$ is constructible in U (so also in X). The Boolean algebra of locally constructible subsets of a locally small space may be strictly larger than the Boolean algebra of constructible subsets (to see this one can construct a sequence X_n of constructible subsets of some small spaces Z_n each X_n needing at least n open sets in the description, and then glue the spaces Z_n into one locally small space).

We will say that a locally small space has the **closure property** if the following holds:

(CP) the weak closure of a small locally closed subset is a closed subset.

Fact 4.11. *Each locally small space with a regular strong topology and with the closure property is weakly regular.*

Fact 4.12. *Each locally constructible subset of a locally small space is a locally essentially finite union of small locally closed subsets.*

Theorem 4.13. *If a locally small space has the closure property, then the weak closure of each locally constructible set is a closed set.*

Proof. By the Fact 4.12, the weak closure of each locally constructible set is a union of weak closures of some locally essentially finite family of small constructible sets. But the family of these closures is also locally essentially finite, and the thesis follows from Theorem 4.4. $\square$

By the above, if a locally small space has the closure property, then the closure operator of the generated topology restricted to the class of locally constructible sets may be considered as the *closure operator* of the generalized topology (in general, the closure operator on a *gts* does not exist).

Notice that each subset Y of a locally small space X forms a (locally small) subspace, since if $X = \bigcup_{\alpha}^a X_{\alpha}$ with all X_{α} small open, then the formula $Y = \bigcup_{\alpha}^a (X_{\alpha} \cap Y)$ defines a locally small space that is a subspace of X . Hence Proposition 3.17, the concept of a connected component, and a quasi-component extend to the category **LSS**.

5 Weakly small spaces

A **weakly** (or **piecewise**) **small space** is a *gts* X having a family $(X_{\alpha})_{\alpha \in A}$ of closed small subspaces indexed by a partially ordered set A such that the following conditions hold:

- W1) X is the union of all X_{α} 's as sets,
- W2) if $\alpha \leq \beta$ then X_{α} is a (closed, small) subspace of X_{β} ,
- W3) for each $\alpha \in A$ there are only finitely many $\beta \in A$ such that $\beta < \alpha$,
- W4) for each two $\alpha, \beta \in A$ there is $\gamma \in A$ such that $X_{\alpha} \cap X_{\beta} = X_{\gamma}$,
- W5) for each two $\alpha, \beta \in A$ there is $\gamma \in A$ such that $\gamma \geq \alpha$ and $\gamma \geq \beta$,
- W6) the *gts* X is the inductive limit of the directed family $(X_{\alpha})_{\alpha \in A}$, which means:
 - a) a subset U of X is open iff all sets $X_{\alpha} \cap U$ are open in respective X_{α} s,
 - b) an open family $\mathcal{U}$ is admissible iff all for each $\alpha \in A$ the family $\mathcal{U} \sqcap X_{\alpha}$ is admissible (=essentially finite) in respective X_{α} .

Such a family $(X_{\alpha})_{\alpha \in A}$ is called an **exhaustion** of X . The weakly small spaces form a full subcategory **WSS** of **GTS**. Here “admissible” means “piecewise essentially finite”, where “piecewise” means “when restricted to a member of the exhaustion (chosen to witness that X is a weakly small space)”. Members of this exhaustion of X may be called **pieces**. If $(X_{\alpha})_{\alpha \in A}$ is an exhaustion of X , then we will write $X = \bigcup_{\alpha \in A}^e X_{\alpha}$.

The **index function** for the exhaustion $(X_{\alpha})_{\alpha \in A}$ is the function $\eta : X \rightarrow A$ given by the formula

$$\eta(x) = \inf\{\alpha \in A \mid x \in X_{\alpha}\}.$$

Here infimum exists thanks to W3). The index function η gives a decomposition of the space X into small locally closed subspaces $X_{\alpha}^0 = \eta^{-1}(\alpha) = X_{\alpha} \setminus \bigcup_{\beta < \alpha} X_{\beta}$.

Fact 5.1. *A subset Y of a weakly small space X is closed if and only if it is piecewise closed.*

Proposition 5.2. *A piecewise essentially finite union of closed subsets of a weakly small space is closed.*

Proof. An essentially finite union of closed subsets is closed. Hence a piecewise essentially finite union of closed sets is piecewise closed, thus closed. $\square$

Notice that the chosen exhaustion of a weakly small space is a piecewise essentially finite (relative to this exhaustion) family of closed sets, and remind that a constructible subset of a piece is a finite union of locally closed subsets of a piece. A **weakly** (or **piecewise**) **constructible subset** is such a subset $Y \subseteq X$ that has constructible intersections with all members of the chosen exhaustion $(X_{\alpha})_{\alpha \in A}$.

Proposition 5.3. *Piecewise constructible subsets of a weakly small space are exactly piecewise essentially finite unions of locally closed subsets of pieces.*

Proof. Piecewise constructible subsets are piecewise essentially finite unions of locally closed subsets of pieces. An essentially finite union of locally closed subsets of pieces is a finite union of locally closed subsets of a single piece, so a constructible subset of a piece. Now apply “piecewise”. $\square$

The **strong topology** on $X = \bigcup_{\alpha \in A}^e X_\alpha$ is the topology that makes the topological space X the respective inductive limit of the system of topological spaces X_α . Its members are all piecewise weakly open subsets, not only weakly (piecewise) open subsets. Hence the open sets from the generalized topology may not form a basis of the strong topology (see Appendix C of [6]). Another unpleasant fact about the weakly small spaces (comparing with the locally small spaces) is that points may not have small neighborhoods (consider an infinite wedge of circles as in Example 4.1.8 of [6]).

We will say that a weakly small space has the **closure property** if the following holds:

(**CP**) the weak closure of a locally closed subset of a piece is a closed subset.

Notice that the strong topology and the generated topology coincide on every piece. If the space X has the closure property, then the topological closure operator restricted to the class of constructible subsets of pieces of X may be treated as the *closure operator* of the generalized topology.

Notice that each subset Y of a weakly small space X forms a (weakly small) subspace, since if $X = \bigcup_{\alpha}^e X_\alpha$ with all X_α small closed, then $Y = \bigcup_{\alpha}^e (X_\alpha \cap Y)$ defines a weakly small space that is a subspace of X . Hence Proposition 3.17, the concept of a connected component, and a quasi-component extend to the category **WSS**.

Theorem 5.4 (cf. IV.2.1 in [6]). *If a weakly small space $X = \bigcup_{\alpha \in A}^e X_\alpha$ is strongly T_1 , and L is a small space, then for each strictly continuous mapping $f : L \rightarrow X$ there is $\alpha_0 \in A$ such that $f(L) \subseteq X_{\alpha_0}$.*

Proof. Let $\eta : X \rightarrow A$ denote the index function for the exhaustion $(X_\alpha)_{\alpha \in A}$. If $\eta(f(L))$ were infinite, then for each $\alpha \in \eta(f(L))$ we could choose $x_\alpha \in f(L)$ with $\eta(x) = \alpha$ and some $y_\alpha \in f^{-1}(x_\alpha)$. Set $S = \{x_\alpha : \alpha \in \eta(f(L))\}$. For each $\gamma \in A$, the set $S \cap X_\gamma$ is finite. Since X is strongly T_1 , the set S is closed as well as each of its subsets, so S is topological discrete. Then $\{y_\alpha : \alpha \in \eta(f(L))\} \subseteq f^{-1}(S)$ is a topological discrete, infinite, and small subset of L . This is a contradiction.

Hence $\eta(f(L))$ is finite, and there is $\alpha_0 \geq \eta(f(L))$. We get $f(L) \subseteq X_{\alpha_0}$. $\square$

Theorem 5.5 (cf. IV.2.2 in [6]). *If a weakly small space X with an exhaustion $(X_\alpha)_{\alpha \in A}$ is strongly T_1 , then each small subspace L of X is contained in some X_{α_0} . In particular, each member X_β of any exhaustion $(X_\beta)_{\beta \in B}$ is contained in some member X_{α_0} of the initial exhaustion.*

Proof. For each such L , the inclusion mapping $i : L \rightarrow X$ is strictly continuous. By Theorem 5.4, the set $i(L) = L$ is contained in a member of the exhaustion $(X_\alpha)_{\alpha \in A}$. $\square$

Let us denote by **WSS**₁ the full subcategory of **WSS** composed of strongly T_1 objects of **WSS**. In this category the term “piecewise” does not depend on an exhaustion (as it

may be expressed by “when restricted to a closed small set”), hence passing to the strong topology gives a functor $(\)_{stop} : \mathbf{WSS}_1 \rightarrow \mathbf{Top}$.

Theorem 5.6. *The category $\mathbf{WSS}_1$ has finite products.*

Proof. It is enough to consider binary products. If X, Y have exhaustions $(X_\alpha)_{\alpha \in A}, (Y_\beta)_{\beta \in B}$, respectively, then $(X_\alpha \times Y_\beta)_{(\alpha, \beta) \in A \times B}$ is an exhaustion defining the weakly small space $X \times Y$. Notice that $X \times Y$ is strongly T_1 . The projections are obviously strictly continuous, and for strictly continuous $f : Z \rightarrow X, g : Z \rightarrow Y$, the mapping $(f, g) : Z \rightarrow X \times Y$ is strictly continuous, since if $Z = \bigcup_{\gamma \in \Gamma}^e Z_\gamma$, then for each Z_{γ_0} there are X_{α_0} and Y_{β_0} such that $f(Z_{\gamma_0}) \subseteq X_{\alpha_0}$ and $g(Z_{\gamma_0}) \subseteq Y_{\beta_0}$. Thus it is the product of X and Y in $\mathbf{WSS}_1$. $\square$

Fact 5.7. *If X is a small space and Y is an object of $\mathbf{WSS}_1$, then the projection $X \times Y \rightarrow Y$ is an open and closed mapping.*

6 Spaces over structures

Assume that $\mathcal{M}$ is any model-theoretic structure.

A **function sheaf over $\mathcal{M}$** on a *gts* X is a sheaf F of sets on X (the sheaf property is assumed only for admissible coverings) such that for each open U the set $F(U)$ is contained in the set M^U of all functions from U into M , and the restrictions of the sheaf are the set-theoretical restrictions of functions.

A **space over $\mathcal{M}$** is a pair (X, O_X) , where X is a *gts* and O_X is a function sheaf over $\mathcal{M}$ on X . A **morphism** $f : (X, O_X) \rightarrow (Y, O_Y)$ of spaces over $\mathcal{M}$ is a strictly continuous mapping $f : X \rightarrow Y$ such that for each open subset V of Y the set-theoretical substitution $h \mapsto h \circ f$ gives the mapping $f_V^\# : O_Y(V) \rightarrow O_X(f^{-1}(V))$. (We could informally say that $f^\# : O_Y \rightarrow O_X$ is the “morphism of function sheaves” over $\mathcal{M}$ induced set-theoretically by f .) An **isomorphism** is an invertible morphism. We get a category $\mathbf{Space}(\mathcal{M})$ of spaces over $\mathcal{M}$ and their morphisms.

Theorem 6.1. *For each structure $\mathcal{M}$, the category $\mathbf{Space}(\mathcal{M})$ has inductive limits.*

Proof. Assume that X is the inductive limit (in the category $\mathbf{GTS}$) of a diagram D with objects $(X_i)_{i \in I}$, indexed by a small category I , and the canonical morphisms $\phi_i : X_i \rightarrow X$.

Assume additionally that function sheaves O_{X_i} over $\mathcal{M}$ are given. Then define

$$O_X(U) = \{h : U \rightarrow M \mid \forall i \in I \ h \circ \phi_i \in O_{X_i}(\phi_i^{-1}(U))\}.$$

Since all O_{X_i} are function sheaves and the section amalgamation in function sheaves is given by the union of graphs of sections, also O_X is a function sheaf. By the above definition, each ϕ_i is a morphism of spaces over $\mathcal{M}$.

If another space (Y, O_Y) with a set of morphisms $\psi_i : X_i \rightarrow Y$ is given, then there is a unique morphism $\eta : X \rightarrow Y$ in $\mathbf{GTS}$ such that $\psi_i = \eta \circ \phi_i$ for all i . But η is also a morphism in $\mathbf{Space}(\mathcal{M})$, because the condition $k \circ \eta \in O_X(\eta^{-1}(W))$ for each $k \in O_Y(W)$ is visibly satisfied, since, for each $i \in I$, we have

$$k \circ \eta \circ \phi_i = k \circ \psi_i \in O_{X_i}(\psi_i^{-1}(W)) = O_{X_i}(\phi_i^{-1}(\eta^{-1}(W))).$$

$\square$

Now assume that a topology is given on the underlying set M of the structure $\mathcal{M}$. We explicitly demand that the product topologies on cartesian powers M^n should be considered. We will call such $\mathcal{M}$ a **weakly topological structure**. (This setting seems to coincide with the case (i) in the introduction of [11]. We do not explore a special language L_t for topological structures considered in the case (ii) of the introduction of [11] or in [4].) Then all projections $\pi_{n,i} : M^n \rightarrow M^{n-1}$ (forgetting the i -th coordinate) are continuous and open. Thus, for example, the field of complex numbers $(\mathbb{C}, +, \cdot)$ considered with the euclidean topology (but not with the Zariski topology) is a weakly topological structure.

Now for each definable set $D \subseteq M^n$, we set:

- a) open subset of a gts = relatively open, definable subset;
- b) admissible covering of a gts = essentially finite covering.

Each such D becomes a small gts , and $\pi_{n,i}$ become strictly continuous and open. We define the structure sheaf O_D as the sheaf of all definable continuous functions from respective (gts -)open subsets $U \subseteq D$ into M .

An **open subspace** of a space over $\mathcal{M}$ is an open subset of its gts together with the function sheaf of the space restricted to this open set. A **small subspace** K of a space X over $\mathcal{M}$ is a small subset of its gts with the function sheaf O_K composed of all finite open unions of restrictions of sections (of the function sheaf O_X) to the relatively open subsets of the small subspace K .

An **affine definable space** over $\mathcal{M}$ is a space over $\mathcal{M}$ isomorphic to a definable subset of some M^n . A **definable space** over $\mathcal{M}$ is a space over $\mathcal{M}$ that has a finite open covering by affine definable subspaces. (The structure sheaf is determined in an obvious way.) A **locally definable space** over $\mathcal{M}$ is a space over $\mathcal{M}$ that has an admissible covering by affine definable open subspaces. (The sections of a structure sheaf are admissible unions of sections of the structure sheaves of affine open subspaces.) A **weakly definable space** X over $\mathcal{M}$ is a space over $\mathcal{M}$ that has an exhaustion $(X_\alpha)_{\alpha \in A}$ composed of definable (small) subspaces such that a function $h : V \rightarrow M$ is a section of O_X iff all restrictions $h|_{V \cap X_\alpha}$ are sections of respective O_{X_α} s.

Morphisms of affine definable spaces, definable spaces, locally definable spaces, and weakly definable spaces over $\mathcal{M}$ are their morphisms as spaces over $\mathcal{M}$. We get full subcategories **ADS**($\mathcal{M}$), **DS**($\mathcal{M}$), **LDS**($\mathcal{M}$), **WDS**($\mathcal{M}$) of **Space**($\mathcal{M}$).

Notice that each definable subset of an affine definable space is also an affine definable space. Thus definable subsets of definable spaces are naturally **subspaces** in **DS**($\mathcal{M}$). Locally definable subsets of locally definable spaces are naturally **subspaces** in **LDS**($\mathcal{M}$). Also piecewise definable subsets of weakly definable spaces are naturally **subspaces** in **WDS**($\mathcal{M}$).

The notions of paracompactness and Lindelöfness for locally definable spaces coincide with their counterparts for the underlying locally small spaces.

Just from the definitions, we get

Facts 6.2. *Definable spaces are small. Every small subspace of a locally definable space is definable.*

Notice that if M is T_1 , then all objects of **LDS**($\mathcal{M}$) and **WDS**($\mathcal{M}$) are strongly T_1 . Moreover, each weakly T_1 locally definable or weakly definable space over $\mathcal{M}$ is strongly T_1 . (It is visible for affine spaces, and extends to the general case by applying “locally”

or “piecewise”.) Thus we can speak just about T_1 objects of $\mathbf{LDS}(\mathcal{M})$ or of $\mathbf{WDS}(\mathcal{M})$ for any $\mathcal{M}$. From Theorem 5.4, we get

Fact 6.3. *If a weakly definable space is T_1 , then each small subspace is contained in a piece, hence it is a definable subspace.*

Theorem 6.4. *For each weakly topological structure $\mathcal{M}$, the products of small spaces in the category $\mathbf{Space}(\mathcal{M})$ exist.*

Proof. Let a family $\{(X_\alpha, O_{X_\alpha})\}_{\alpha \in A}$ of small spaces over $\mathcal{M}$ be given. By the Proposition 3.11, the family $\{X_\alpha\}_{\alpha \in A}$ has the product X in $\mathbf{GTS}$. Now, for any $h \in O_{X_\alpha}(V)$, the function $h \circ \pi_\alpha : \pi_\alpha^{-1}(V) \rightarrow M$ may be called a “function of one variable”, where $\pi_\alpha : X \rightarrow X_\alpha$ is the projection. Define sections of O_X as all admissible (so essentially finite) unions of compatible functions of one variable. Then the projections π_α are morphisms of $\mathbf{Space}(\mathcal{M})$. For a family $f_\alpha : Y \rightarrow X_\alpha$ of morphisms of $\mathbf{Space}(\mathcal{M})$, the induced mapping $(f_\alpha)_{\alpha \in A} : Y \rightarrow X$ is a morphism of $\mathbf{Space}(\mathcal{M})$, since all the sections of O_X are finite unions of the functions of one variable, which are determined by sections of O_{X_α} s. $\square$

Unfortunately, even finite products of affine definable spaces usually are not affine definable spaces.

Theorem 6.5. *For each weakly topological structure $\mathcal{M}$, finite products in $\mathbf{ADS}(\mathcal{M})$ exist.*

Proof. For definable sets $D \subseteq M^n$ and $E \subseteq M^m$, the product is the definable set $D \times E \subseteq M^{n+m}$ (with its natural space structure). Indeed, the projections are continuous definable mappings, and for any continuous definable $f : Z \rightarrow D$, $g : Z \rightarrow E$, with Z definable in some M^k , the induced mapping $(f, g) : Z \rightarrow D \times E$ is continuous definable, since the topology of $D \times E$ is the product topology. For a continuous definable function $h \in O_{D \times E}(U)$, the function $h \circ (f, g) : (f, g)^{-1}(U) \rightarrow M$ is continuous definable. This proves finite products exist in $\mathbf{ADS}(\mathcal{M})$. $\square$

Notice that the canonical projections from a finite product of affine definable spaces to its factors are open morphisms, but not in general closed morphisms.

Fact 6.6 (cf. I.1.3 in [3]). *A mapping $f : X \rightarrow Y$ between objects of $\mathbf{LDS}(\mathcal{M})$ is a morphism iff the image of each open definable subspace is contained in an open definable subspace and when restricted to open definable subspaces in the domain and in the range, the mapping is continuous definable.*

Theorem 6.7 (cf. I.2.5 in [3]). *For each weakly topological structure $\mathcal{M}$, the category $\mathbf{LDS}(\mathcal{M})$ has finite products.*

Proof. For locally definable spaces $(X, O_X) = \bigcup_{\alpha \in A}^a (X_\alpha, O_{X_\alpha})$, $(Y, O_Y) = \bigcup_{\beta \in B}^a (Y_\beta, O_{Y_\beta})$, where (X_α, O_{X_α}) , (Y_β, O_{Y_β}) are affine definable subspaces over $\mathcal{M}$, consider the space $\bigcup_{(\alpha, \beta) \in A \times B}^a (X_\alpha, O_{X_\alpha}) \times (Y_\beta, O_{Y_\beta}) = \bigcup_{(\alpha, \beta) \in A \times B}^a (X_\alpha \times Y_\beta, O_{X_\alpha \times Y_\beta})$. This is the product of given locally definable spaces, since the projections are visibly morphisms in $\mathbf{LDS}(\mathcal{M})$, and for given morphisms $f : Z \rightarrow X$, $g : Z \rightarrow Y$ in $\mathbf{LDS}(\mathcal{M})$ from the space $(Z, O_Z) = \bigcup_{\gamma \in \Gamma}^a (Z_\gamma, O_{Z_\gamma})$, the induced mapping $(f, g) : Z \rightarrow X \times Y$ is a morphism

in $\mathbf{LDS}(\mathcal{M})$. Indeed, we may assume $\{Z_\gamma\}_{\gamma \in \Gamma}$ is an admissible covering by affine definable spaces that is a refinement of the preimage of the covering $\{X_\alpha \times Y_\beta\}_{(\alpha, \beta) \in A \times B}$ and thus each Z_γ is mapped into some $X_\alpha \times Y_\beta$ by some continuous definable mapping, and the whole (f, g) is an admissible union of such partial mappings. $\square$

A subset Y of a locally definable space X (over $\mathcal{M}$) may be called **local** if for each $y \in Y$ there is an open small neighborhood U of y such that $Y \cap U$ is definable in U . On such Y , we could define a locally definable space by the formula

$$Y = \bigcup^a \{Y \cap U \mid U \text{ small, open, } Y \cap U \text{ definable in } U\}.$$

(This definition is canonical in the sense that it does not depend on any arbitrary choice of open neighborhoods.) However, the use of local subsets does not reflect this locally definable space (see Example 10 in [10]). Any weakly open and any weakly discrete set is local.

Let us denote by $\mathbf{WDS}_1(\mathcal{M})$ the full subcategory of T_1 objects of $\mathbf{WDS}(\mathcal{M})$ for any $\mathcal{M}$.

Fact 6.8 (cf. IV.2.3 in [6]). *A mapping $f : X \rightarrow Y$ between objects of $\mathbf{WDS}_1(\mathcal{M})$ is a morphism iff the image of a piece is contained in a piece and when restricted to pieces in the domain and in the range, the mapping is continuous definable.*

Notice that for two morphisms $f_1 : X_1 \rightarrow Y_1$, $f_2 : X_2 \rightarrow Y_2$ of $\mathbf{LDS}(\mathcal{M})$ or of $\mathbf{WDS}_1(\mathcal{M})$, their cartesian product $f_1 \times f_2 : X_1 \times X_2 \rightarrow Y_1 \times Y_2$ is a morphism.

Theorem 6.9. *For each weakly topological structure $\mathcal{M}$, the category $\mathbf{WDS}_1(\mathcal{M})$ has finite products.*

Proof. Notice that by Theorem 6.7, finite products exist in the category $\mathbf{DS}(\mathcal{M})$, and the product of T_1 spaces is T_1 . We will drop the structure sheaves in notation. If $(X_\alpha)_{\alpha \in A}$, $(Y_\beta)_{\beta \in B}$ are exhaustions of objects X , Y , then define $X \times Y = \bigcup_{(\alpha, \beta) \in A \times B}^e (X_\alpha \times Y_\beta)_{(\alpha, \beta) \in A \times B}$. This is an exhaustion determining the structure sheaf. The resulting space is clearly T_1 , and the projections $X \times Y \rightarrow X$, $X \times Y \rightarrow Y$ are visibly morphisms of $\mathbf{WDS}_1(\mathcal{M})$. If $Z = \bigcup_{\gamma \in \Gamma}^e Z_\gamma$ is any object of $\mathbf{WDS}_1(\mathcal{M})$ and $f : Z \rightarrow X$, $g : Z \rightarrow Y$ are morphisms, then the mapping $(f, g) : Z \rightarrow X \times Y$ is a morphism because: we may assume that for each Z_{γ_0} there are X_{α_0} and Y_{β_0} such that $f(Z_{\gamma_0}) \subseteq X_{\alpha_0}$ and $g(Z_{\gamma_0}) \subseteq Y_{\beta_0}$, the generalized topology of the product of definable spaces contains the generalized topology of their (generalized topological) product, and the definable continuous mappings are stable under composition and juxtaposition. Thus $X \times Y$ is the product of X and Y in $\mathbf{WDS}_1(\mathcal{M})$. $\square$

Theorem 6.10. *For any weakly topological structure $\mathcal{M}$, the category $\mathbf{LDS}(\mathcal{M})$ has fiber products.*

Proof. The fiber product of morphisms $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ is the subspace $(f \times g)^{-1}(\Delta_Z) \subseteq X \times Y$, where Δ_Z is the diagonal of the space Z (locally definable in $Z \times Z$). $\square$

Theorem 6.11. *For any weakly topological structure $\mathcal{M}$, the category $\mathbf{WDS}_1(\mathcal{M})$ has fiber products.*

The proof of this theorem is similar to the proof of the previous theorem.

Let $\mathcal{C}$ be one of the categories: $\mathbf{LDS}(\mathcal{M})$ or $\mathbf{WDS}_1(\mathcal{M})$. We will say that an object Z of $\mathcal{C}$ is **$\mathcal{C}$ -complete** if the mapping $Z \rightarrow \{*\}$ is universally closed in $\mathcal{C}$, which means that for each object Y of $\mathcal{C}$ the projection $Z \times Y \rightarrow Y$ (which is the base extension of $Z \rightarrow \{*\}$) is a closed mapping.

Proposition 6.12. *A closed subspace of a $\mathcal{C}$ -complete space is $\mathcal{C}$ -complete.*

Proof. Let C be a closed subspace of a $\mathcal{C}$ -complete space Z . For any Y in $\mathcal{C}$, the space $C \times Y$ is a closed subset of $Z \times Y$. Thus the image under the projection along C of any closed subset of $C \times Y$ is closed in Y . $\square$

Proposition 6.13. *Every subspace of an object of $\mathcal{C}$ that is closed in the generated topology is a closed subspace.*

Proof. This is clear for affine definable spaces. In general, closedness may be checked “locally” or “piecewise”. $\square$

Proposition 6.14. *For an object Z of $\mathcal{C}$, the following conditions are equivalent:*

- a) Z is weakly Hausdorff;
- b) Z is strongly Hausdorff;
- c) Z has its diagonal Δ_Z closed.

Proof. If a space X is weakly Hausdorff, then its generated topology is Hausdorff, thus the diagonal Δ_X is closed in the generated topology. But Δ_X is always a subspace, thus, by Proposition 6.13, a closed subspace.

If the diagonal Δ_X of a space X is a closed subspace, then the generated topology of X is Hausdorff, so X is weakly Hausdorff. But it is also strongly T_1 , since for any $x_0 \in X$, the set $\{(x_0, x_0)\}$ is closed in $\{x_0\} \times X$ and the projection $\{x_0\} \times X \rightarrow X$ is closed. Thus X is strongly Hausdorff. $\square$

Notice that if the topology of M is Hausdorff, then:

- a) all M^n as well as all objects of $\mathbf{ADS}(\mathcal{M})$ are (strongly) Hausdorff;
- b) all objects of $\mathbf{WDS}(\mathcal{M})$ having exhaustions consisting of affine definable spaces are (strongly) Hausdorff;
- c) an object of $\mathbf{DS}(\mathcal{M})$ may not be (weakly) Hausdorff.

Because of 6.14, we may speak just about Hausdorff objects of $\mathcal{C}$.

Proposition 6.15. *If Y is a Hausdorff object of $\mathcal{C}$, then:*

- a) *each of its $\mathcal{C}$ -complete subspaces is closed;*
- b) *the graph of each morphism $f : X \rightarrow Y$ in $\mathcal{C}$ is closed;*
- c) *the image of a $\mathcal{C}$ -complete subspace C under a morphism $g : Y \rightarrow Z$ in $\mathcal{C}$, with Z Hausdorff, is closed and complete.*

Proof. a) If C is a $\mathcal{C}$ -complete subspace of Y , then $\Delta_Y \cap (C \times Y)$ is relatively closed in $C \times Y$, and its projection on Y , equal to C , is closed.

b) The graph of f is a subspace of $X \times Y$ being the preimage of Δ_Y by a morphism $f \times id_Y$.

c) The graph of $g|_C$ is closed in $C \times Z$. Since C is complete, the projection along C of the graph, that is $g(C)$, is closed. Now, in order to prove that $g(C)$ is complete, consider an object W of $\mathcal{C}$. Then for a closed subset A of $g(C) \times W$, we have $\pi_{g(C)}(A) = \pi_C((g|_C \times id_W)^{-1}(A))$ is a closed set. $\square$

Suppose $\mathcal{M}$ satisfies the condition
 (DCCD) $\mathcal{M}$ admits (finite) cell decomposition with definably connected cells
 (here we assume only that a cell is a definable set, for stronger notions of a cell see [9] or [12]), then:

- a) each object of $\mathbf{DS}(\mathcal{M})$ has a finite number of clopen definable connected components, and is a finite (generalized) topological direct sum of them;
- b) each object of $\mathbf{LDS}(\mathcal{M})$ is a (locally finite) direct sum of its connected components;
- c) each object of $\mathbf{WDS}(\mathcal{M})$ is a piecewise finite union of its clopen connected components, so also a direct sum of its connected components.

Thus If $\mathcal{M}$ satisfies (DCCD), then all objects of $\mathbf{LDS}(\mathcal{M})$ and of $\mathbf{WDS}(\mathcal{M})$ satisfy (ACC). All o-minimal structures $\mathcal{M}$ with their order dense satisfy (DCCD), but t-minimality in the sense of [12] does not guarantee (DCCD).

Theorem 6.16. *If $\mathcal{M}$ satisfies (DCCD) and there is a Hausdorff object Z of $\mathcal{C}$ having a non-closed countable subset, then each $\mathcal{C}$ -complete space has a finite number of connected components.*

Proof. The $\mathcal{C}$ -complete space C is a direct sum of the family of its connected components. Take one point from each connected component. The resulting subspace S is discrete and closed, so complete by Proposition 6.12. If it were infinite, then we could assume (by taking a closed subspace if necessary) it is countable. An injection from S to a countable non-closed subset of some Hausdorff object of $\mathcal{C}$ would give a counterexample of c) in Proposition 6.15. Hence S is finite. $\square$

Theorem 6.17. *Assume that $\mathcal{M}$ satisfies (DCCD) and there is a Hausdorff object Z of $\mathcal{C}$ having a non-closed countable subset. Let C be a $\mathcal{C}$ -complete space, where if $\mathcal{C} = \mathbf{LDS}(\mathcal{M})$ then C is assumed to be T_1 and have Lindelöf connected components. Then C is a definable space.*

Proof. In the case of a locally definable space: we can assume by Theorem 6.16 that C is connected. If $C = \bigcup_{n \in \mathbb{N}}^a C_n$ is an admissible covering by affine definable spaces and C is not definable, then we may assume for each $n \in \mathbb{N}$ we can choose $x_n \in C_n \setminus (C_0 \cup \dots \cup C_{n-1}) \neq \emptyset$. The set $B = \{x_n | n \in \mathbb{N}\}$ is infinite, not small, locally finite, so, by T_1 , topological discrete. Moreover, B is closed, thus complete (by Proposition 6.12).

In the case of a weakly definable space: if C is not definable, then the index function η of an exhaustion $(C_\alpha)_{\alpha \in A}$ has infinite image, and we can choose an element $x_\alpha \in C_\alpha^0$ for each $\alpha \in \eta(C)$. The set $B = \{x_\alpha : \alpha \in \eta(C)\}$ is an infinite (but piecewise finite) closed subspace all of whose subsets are also closed. Thus B is a complete (by Proposition 6.12) and topological discrete space. We may assume B is countable.

In both cases an injection $\phi : B \rightarrow Z$ with the image a countable non-closed set would be a morphism giving contradiction with c) of Proposition 6.15. Thus a countable set B cannot be constructed and C is definable. $\square$

Remind (after [11]) that a structure $\mathcal{M}$ is a **first order topological structure** (called in [4] a topological structure with explicitly definable topology) if the basis of the topology on M is uniformly definable in $\mathcal{M}$. (There is a formula $\Phi(x, \bar{y})$ of the (first order) language of $\mathcal{M}$ such that the family $\{\Phi(x, \bar{a})^M : \bar{a} \subseteq M\}$ is the basis of the topology of M .) If $\mathcal{M}$ is a first order topological structure, then the relative closure of a definable subset in a definable set is definable (see [11]). Thus the closure operator

for the generalized topology exists for the locally definable subsets of locally definable spaces, and for definable subsets of pieces in weakly definable spaces over $\mathcal{M}$.

Acknowledgements. This paper was written during my stay at the Fields Institute during the Thematic Program on o-minimal Structures and Real Analytic Geometry in 2009. I thank the Fields Institute for their warm hospitality.

References

- [1] E. Baro, M. Otero, *Locally definable homotopy*, Annals of Pure and Applied Logic (2009), doi:10.1016/j.apal.2009.03.003.
- [2] M. Barr, C. Wells, *Toposes, Triples and Theories*, Grundlehren der mathematischen Wissenschaft 278, Springer-Verlag 1985.
- [3] H. Delfs, M. Knebusch, *Locally Semialgebraic Spaces*, Lecture Notes in Mathematics 1173, Springer-Verlag 1985.
- [4] J. Flum, M. Ziegler, *Topological Model Theory*, Lecture Notes in Mathematics 769, Springer-Verlag 1980.
- [5] M. Kashiwara, P. Shapira, *Categories and Sheaves*, Grundlehren der mathematischen Wissenschaften 332, Springer-Verlag 2006.
- [6] M. Knebusch, *Weakly Semialgebraic Spaces*, Lecture Notes in Mathematics 1367, Springer-Verlag 1989.
- [7] M. Knebusch, *Semialgebraic topology in the recent ten years*, "Real Algebraic Geometry Proceedings, Rennes 1991", M. Coste, L. Mahé, M.-F. Roy, eds., LNM 1524, Springer 1992, 1–36.
- [8] S. Mac Lane, I. Moerdijk, *Sheaves in Geometry and Logic*, Universitext, Springer-Verlag 1992.
- [9] Mathews, *Cell decomposition and dimension functions in first order topological structures*, Proc. London Math. Soc. (3) 70 (1995), 1–32.
- [10] A. Piękosz, *O-minimal homotopy and generalized (co)homology*, arXiv:0808.3866 [math.LO].
- [11] A. Pillay, *First order topological structures and theories*, Journal of Symbolic Logic Vol. 53, No. 3 (1987), 763–778.
- [12] H. Schoutens, *T-minimality*, preprint, 2001.

POLITECHNIKA KRAKOWSKA
 INSTYTUT MATEMATYKI
 WARSZAWSKA 24
 PL-31-155 KRAKÓW
 POLAND
 E-mail: pupiekos@cyf-kr.edu.pl